

## **EFFECT OF SULFUR AND APPLICATION STAGES ON YIELD AND YIELD COMPONENTS OF SOYBEAN**

**Salman Khan**

The University of Agriculture  
Peshawar-Pakistan

**Muhammad Asim**

The University of Agriculture  
Peshawar-Pakistan

**Muhammad Usaid**

The University of Agriculture  
Peshawar-Pakistan

**Hamza Masud**

The University of Agriculture  
Peshawar-Pakistan

**Abdul Wahab**

Directorate of soil and plant  
nutrition ARI Tarnab Peshawar-  
Pakistan

**Abdul Haseeb**

The University of Agriculture  
Peshawar-Pakistan.

**Asif Ismail**

The University of Agriculture  
Faisalabad-Pakistan

**Hazrat Ali**

Department of health and  
biological sciences, Abasyn  
University Peshawar-Pakistan

**Rizwan Ali**

The University of Agriculture  
Peshawar-Pakistan

**Uzair Ahmed**

The University of Agriculture  
Peshawar-Pakistan

**Sajid Ali**

The University of Agriculture  
Peshawar-Pakistan.

### Article Info



### Abstract

*Soybean is a vital crop for food and oil production, yet its yield and quality are often compromised by suboptimal nutrient management. This study aimed to address this issue by investigating the effects of sulfur and its time of application on soybean at the Research Farm of The University of Agriculture Peshawar during Kharif season of 2022. The experiment followed a randomized complete block design with three replicates. It aimed to evaluate the effects of four sulfur levels (0, 20, 40, and 60 kg ha<sup>-1</sup>) applied at three different growth stages (S1=sowing, V3=vegetative, and R1=flowering) using elemental sulfur as the sulfur source. The results showed significant impacts, particularly when sulfur was applied at a rate of 40 kg ha<sup>-1</sup>. This treatment extended days to flowering (56), improved yield components as well as biological (7917 kg ha<sup>-1</sup>) and seed yield (2882 kg ha<sup>-1</sup>). Sulfur at 40 kg ha<sup>-1</sup> positively influenced oil content (18.95%), elevated oil yield (571 kg ha<sup>-1</sup>) and protein content (32.39%). In contrast supplementation of sulfur at 60 kg ha<sup>-1</sup> extended days to maturity and improved the harvest index. Regarding application stages sulfur applied at V3 stage delayed flowering (56), increased leaves count and plant height. Application at R1 stage increased days to maturity (129), yield components, biological and seed yield (7992 and 2806 kg ha<sup>-1</sup>, respectively), oil content and yield (556 kg ha<sup>-1</sup>, 19.54%, respectively). Moreover, yield and yield components showed positive correlation with quality components of soybean. In last, economic analysis revealed high net benefit and BC ratio by applying 40 kg ha<sup>-1</sup> of sulfur. In conclusion, applying 40 kg ha<sup>-1</sup> of sulfur at the R1 stage is highly effective and economically feasible in enhancing soybean yield and quality under the agro-climatic conditions of Peshawar. Future research should explore the long-term effects of sulfur application on soil health and productivity, as well as the economic feasibility of these practices for large-scale adoption*



This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license <https://creativecommons.org/licenses/by/4.0>

**Keywords:** Growth stages, Oil content, Plant height, Protein content, Peshawar, Soybean yield, Sulfur levels

## Introduction

Soybean is an annual summer crop originated from East Asia and is extensively cultivated for its consumable beans. It contributes significantly to global oilseed production, representing 58 %, and constitutes 69 % of the protein meal consumed by cattle worldwide. Soybean is regarded as one of the paramount protein and oil crops on a global scale. Its cultivation holds immense significance in the agricultural sector. It has proven to be a globally viable industrial and commercial oilseed crop and known as the 'Golden Bean' due to its excellent nutritional content. The crop has generated 9.47 million tons production of soybean after being planted on 8.85 million hectares throughout the world (Berger et al., 2010). Because of its nutritional value, their bi-products such as soya milk, soybean sprouts, and soya nuts, various types of tofu, cottage, cheese, and curd are in high demand. Soybeans are grown all over the world, yielding 276,406,003 tons with over 30% produced in America. Given its remarkable adaptability across diverse industries, soybean holds high economic value on a global scale, estimated at a staggering 119,516 million US dollars. Furthermore, beyond its economic impact, soybean also plays a role in mitigating symptoms associated with certain chronic conditions (Kahraman, 2017). For soybean production to keep up with the expected global population growth rate, several hurdles must be overcome. In Khyber Pakhtunkhwa and Pakistan, annual soybean production is 46 and 66 tons with 54 and 84 hectares of planted area (MNFS&R, 2014) that is not in tandem with its demand.

Sulfur, as an integral component of sulfur-containing amino acids and various metabolites, serves a critical role in various

plant growth and developmental processes. Acknowledged as the 4<sup>th</sup> primary nutrient following nitrogen, phosphorus, and potassium, sulfur's significance in sustaining plant health and growth is widely recognized. Sulfur's significance in soybean seed development has been well documented a number of investigators have reported (Dubey et al., 1995). Sulfur interacts with nitrogen in soybeans, affecting nitrogen metabolism and utilization efficiency, making it vital for proper seed development and yield (Głowacka et al., 2023). Various stages of S application have a substantial impact on the quality, growth, and productivity of canola. Oil, glucosinolate, and protein content are all improved when 20 kg ha<sup>-1</sup> S is applied at plantation (Ahmad et al., 2007). Sulfur application at planting time increased maturity in canola, but there was no change in plant height whether sulfur was treated at the sowing or flowering stage. When S was administered after 42 days of seeding, it resulted in lower seed production on one site, but better seed yield and improved S absorption both in seed and biomass when treated earlier (Karami et al., 2016). Sulfur is required for many plant growth and developmental processes. Plant nutrient deficiencies are the most significant restrictions to crop growth. Sulfur is important element affecting plant growth and yield in the soil (seervi et al., 2018).

Sulfur is essential for numerous plant processes, including photosynthesis, chlorophyll production, and nitrogen fixation and is required for protein synthesis. Cysteine and methionine are two sulfur-containing amino acids found in soybean crops. Soybean uses nearly the same amount of sulfur as they use phosphorus (Paul et al., 2019). Sulfur application can result in increased soybean

yield and dry matter. However, lack of sulfur can result in increased nitrates in soybean seeds (Zhao et al., 2006). When sulfur levels are low, deficiency can occur, causing pale green colour, diminished plant growth and plants require sulfate ion ( $\text{SO}_4$ ), and however, elemental sulfur is still not plant-available and should be transformed to the sulfate form by soil bacteria. Sulfur (Vora et al., 2018). Maintaining an optimal level of sulfur in the soil is pivotal for promoting the production of substantial quantities of high-quality seeds of soybean. Generally, adequate sulfur application significantly enhances both crop growth and the quality of oilseed crops by amplifying protein and oil content (Ahmad et al., 2018). To optimize seed and oil yield, sulfur should be administered in split doses. This approach ensures that the crop receives adequate sulfur throughout its growth and development stages. Applying sulfur in split doses ensures a continuous supply of these essential nutrients to the crop. Notably, applying sulfur before flowering is particularly advantageous as it enhances both the photosynthetic efficiency and the crop's potential yield. It's crucial to note that after the flowering stage, the crop ceases to expand its foliage, limiting further enhancement in yield potential, emphasizing the significance of timely sulfur application before this critical growth stage (Ahmad et al., 2011).

Because the majority of soils in soybean-growing areas are poor in accessible sulfur, and their correct time of application is still in doubt. A systematic study of sulfur nutrition is required in order to increase soybean productivity. Therefore, a trial was performed by subjecting soybean crop to various sulfur levels applied at different application stages to find out the best level and time of application in Peshawar region.

## MATERIALS AND METHODS

The experiment was conducted during the summer of 2022 at the University of Agriculture Peshawar, utilizing a randomized complete block design (RCBD) with three replications. Each plot measured 3 meters by 4 meters, providing a net area of 12  $\text{m}^2$ . Within each plot, six rows were planted, with a spacing of 50 cm between each row. The study involved the commencement of four sulfur levels (0, 20, 40, and 60  $\text{kg ha}^{-1}$ ) at three stages: during sowing, at the vegetative stage (V3), and at the flowering stage (R1). These factors were chosen for ensuring the nutrient is available at critical stages to support optimal development, yield and quality and to evaluate which level and which time of application is the best. Elemental sulfur served as the sulfur source, and the Malakand-96 variety was planted at a rate of 80  $\text{kg ha}^{-1}$ . To fulfill the requirement of micronutrients, N, P and K were applied at 40, 40 and 20  $\text{kg ha}^{-1}$  from urea, DAP and MOP, respectively. Three irrigation sessions were conducted throughout the season: the first three weeks after emergence, the second at the flowering stage, and the third during the pod-filling stage. Manual weeding was performed using a hand hoe, and all necessary cultural practices were consistently applied throughout the trial duration. Following parameters were studied during the trail.

### *Days to flowering and maturity*

It was collected by tallying the days from the initial planting date to the emergence of flowers within each individual plot. Similarly, the maturity days of the crop were documented by tracking the duration from the sowing date to the point at which the pods exhibited a yellowish color within each subplot.

### *Growth attributes*

Plant height (cm) measurements were collected by randomly selecting four plants within each plot and measuring from the base to the top of the plant. These measurements were then averaged to represent the overall plant height for that specific plot. Similarly, the number of leaves per plant was determined by randomly selecting five plants within each plot and counting the number of leaves on each plant. The counts were then averaged to obtain the number of leaves per plant for the entire plot.

#### *Yield and yield attributes*

At maturity, pods on five selected plants from the two central rows were counted and averaged to determine pods per plant. For pod length, ten pods were randomly selected from five plants within each plot and measured, with the average length calculated. To record thousand grain weight (TGW), mature pods were dried, and the weight of 100 grains was multiplied by 10. Biological yield ( $\text{kg ha}^{-1}$ ) was obtained by sun drying and weighing the harvested central two rows. Grain yield ( $\text{kg ha}^{-1}$ ) was determined by drying, threshing, and weighing the yield from the central two rows. The harvest index was calculated by dividing grain yield by biological yield and multiplying by 100.

#### *Soy bean quality*

To determine oil content, a 2-gram dried sample was placed in a paper thimble attached to a Soxhlet extractor. Ether extracted the components over 5 hours, with the residue dried in an oven at 70-80°C for 2 hours, then weighed. Oil content percentage was calculated following AOAC (1990) guidelines. Oil yield ( $\text{kg ha}^{-1}$ ) was obtained by multiplying the oil content percentage by the total seed yield ( $\text{kg ha}^{-1}$ ) from each subplot, then dividing by 100. Protein content of soybean grains was assessed using the Kjeldahl distillation method following sulfuric acid mineralization (Kjeldahl, 1883).

#### *Economic analysis*

Economic yield was obtained by below mentioned equation:

$$\text{Economic yield} = \frac{\text{Total cost}}{\text{Total income}}$$

#### *Statistical analysis*

The collected data were analyzed using analysis of variance (ANOVA) tailored for the randomized complete block (RCB) design. Post hoc comparisons were conducted using the least significant difference test to determine significant differences between treatment means at a probability level of 0.05 (Jan et al., 2009).

*Table 1 Physicochemical properties of experimental soil before plantation at various depths.*

Physicochemical properties of soil	Analyzed values	
Soil depth (cm)	0-20	20-40
Texture	Silt loam	
pH	7.1	7.6
Moisture content (%)	7.41	9.38
Bulk density ( $\text{gcm}^{-3}$ )	1.68	1.71
Porosity (%)	39	36
Organic matter (%)	0.27	0.16
% sand	20.56	20.85
% silt	66.12	65.98
% clay	13.32	13.17
Soil total nitrogen ( $\text{mg kg}^{-1}$ )	513	487
Available P ( $\text{mg kg}^{-1}$ )	14.54	12.79
Available potassium ( $\text{mg kg}^{-1}$ )	82.76	81.89

## RESULTS

Soil analysis of the experimental site was conducted before commencing the field trial to determine soil physicochemical properties. It was observed during the analysis that the soil at the experimental site has a silt loam texture. The recorded values for pH, moisture content, bulk density, porosity, organic matter, soil total nitrogen, available phosphorus, and potassium at depths of 0-20 cm and 20-40 cm are presented in Table 1. Moreover, data on various parameters of soybeans are presented below.

### *Days to flowering*

Significant impacts of sulfur levels and application stages were observed on both growth stages (Table 2). However, the interaction between sulfur levels and application stages was not significant. Delayed flowering, occurring at 56 days, was noted with the use of  $40 \text{ kg S ha}^{-1}$ , while earlier flowering, at 52 days, was observed in the control plots. Similarly, delayed physiological maturity, with 128 days, was recorded in units treated with  $60 \text{ kg ha}^{-1}$  sulfur, whereas control plots exhibited earlier maturity at 125 days. Application of sulfur at the vegetative stage resulted in delayed

flowering with 56 days compared to the sowing stage and control plots. Moreover, plants treated with sulfur at the flowering stage took 129 days to reach maturity compared to those treated at the vegetative

stage, with control plots exhibiting the earliest maturity at 124 days.

*Table 2 Physiology and growth of soybean under different sulfur levels applied at various application stages.*

Application stages (APP)	Days flowering to	Days Maturity to	Plant height (cm)	Leaves plant <sup>-1</sup>
At sowing stage	52 c	124 c	66 b	112 c
At V3 stage	56 a	127 b	68 a	119 a
At R1 stage	55 b	129 a	67 a	115 b
LSD <sub>(0.05)</sub>	1	1	2	2
<b>Sulfur levels (S)</b>				
Control	52 d	125 d	65 c	109 d
20 kg ha <sup>-1</sup>	54 c	126 c	67 b	114 c
40 kg ha <sup>-1</sup>	56 a	127 b	69 a	122 a
60 kg ha <sup>-1</sup>	55 b	128 a	68 ab	117 b
LSD <sub>(0.05)</sub>	1	2	1	3
APP×S	ns	ns	ns	ns

The use of different alphabetic labels within the same groups indicates statistically significant differences at a 5% probability level.

#### *Growth attributes*

Significant variations were observed in growth parameters due to sulfur levels and application timings, although the interaction between these factors was not significant (Table 2). Plants treated with 40 kg ha<sup>-1</sup> sulfur reached maximum height (69 cm), followed closely by those treated with 60 kg ha<sup>-1</sup> sulfur (68 cm), while control plots had the minimum height (65 cm). Similarly, subplots treated with 40 kg ha<sup>-1</sup> sulfur exhibited the highest leaf count (122), with 60 kg ha<sup>-1</sup> sulfur treatments closely behind (117 plant<sup>-1</sup>), whereas control plots had the lowest leaf count (109 plant<sup>-1</sup>). Regarding application stages, plants treated with sulfur at the

vegetative stage (V3) displayed maximum plant height (68 cm) and leaf count (119 plant<sup>-1</sup>), while those treated at the flowering stage (R1) had slightly lower values (67 cm and 115 plant<sup>-1</sup>, respectively).

#### *Yield and yield attributes*

The data analysis revealed significant effects of sulfur levels and application stages on yield and yield attributes of soybean. However, the interactive effect of sulfur rates on various application stages (APP×S) remained non-significant (Table 3). Significant influence of sulfur levels and application stages on pods per plant was observed, with 40 kg S ha<sup>-1</sup> resulting in the highest pod count (69 pods) compared to the control (62 pods). Application at the flowering stage (R1) produced the highest pods per plant (66), followed closely by the vegetative stage (V3) treatment (65 pods),



while the control had the lowest count (64 pods). For pod length, 40 kg ha<sup>-1</sup> sulfur resulting in the longest pods (4.58 cm) compared to the control (3.94 cm). The flowering stage (R1) application also yielded the longest pods (4.50 cm), while the control had the shortest (4.14 cm). TGW varied significantly across different application stages and sulfur levels, with the highest weight observed at the R1 stage (149 g) and with 60 kg ha<sup>-1</sup> sulfur supplementation (149 g), while the control had the lowest TGW (142 g).

As for biological yield, applying 40 kg S ha<sup>-1</sup> resulted in the highest biological yield of 7917 kg ha<sup>-1</sup>, while the control treatment yielded the minimum at 7745 kg ha<sup>-1</sup>. The experimental units treated with 40 kg ha<sup>-1</sup> sulfur exhibited the highest grain yield of 2882 kg ha<sup>-1</sup>, followed by 60 kg ha<sup>-1</sup> sulfur, yielding 2739 kg ha<sup>-1</sup>. Conversely, the control displayed the lowest grain yield of

2547 kg ha<sup>-1</sup>. Moreover, 60 kg ha<sup>-1</sup> sulfur exhibited the maximum harvest Index of 36.78%, followed by 40 kg ha<sup>-1</sup> sulfur, showing a Harvest Index of 34.67%. Conversely, the control plots demonstrated the minimum Harvest Index of 32.92%. Regarding application stages, introducing sulfur at flowering stage exhibited the highest biological yield at 7992 kg ha<sup>-1</sup>, while the lowest yield was recorded when sulfur was applied at the sowing stage, producing 7633 kg ha<sup>-1</sup>. Similarly, for grain yield (kg ha<sup>-1</sup>), the highest yield of 2806 kg ha<sup>-1</sup> was achieved when sulfur application coincided with the flowering stage (R1). Following closely, the treatment during the vegetative stage (V3) resulted in a yield of 2717 kg ha<sup>-1</sup>. Conversely, the lowest yield of 2588 kg ha<sup>-1</sup> was observed with sulfur introduced at the stage of planting. The harvest index of soybean remained unaffected regardless of applying sulfur at various growth stages of soybean (Table 4).

*Table 3 Yield and yield related attributes of soybean under different sulfur levels applied at various application stages.*

Application stages (APP)	Pods plant <sup>-1</sup>	Pod length (cm)	Thousand grain weight (g)	Biological yield (kg ha <sup>-1</sup> )	Grain yield (kg ha <sup>-1</sup> )
At sowing stage	64 b	4.14 b	143 c	7633 c	2588 c
At V3 stage	65 a	4.31 b	146 b	7884 b	2717 b
At R1 stage	66 a	4.50 a	149 a	7992 a	2806 a
LSD <sub>(0.05)</sub>	1	0.2	2	59	117
Sulfur levels (S)					
Control	62 d	3.94 d	142 c	7745 c	2547 c
20 kg ha <sup>-1</sup>	64 c	4.31 c	145 b	7830 b	2645 bc
40 kg ha <sup>-1</sup>	69 a	4.58 ab	147 ab	7917 a	2882 a
60 kg ha <sup>-1</sup>	66 b	4.45 a	149 a	7855 ab	2739 b
LSD <sub>(0.05)</sub>	2	0.2	2	69	101
APP×S	ns	ns	ns	ns	ns

The use of different alphabetic labels within the same groups indicates statistically

significant differences at a 5% probability level.

### Soy bean quality

Mean data regarding quality of soybean showed significant variations when subjected to different sulfur levels at various application stages. The interactive effect (APP×S) remained non-significant except for protein content (Table 4). The highest oil content (18.95%) and oil yield (571 kg ha<sup>-1</sup>) were recorded in plots treated with 40 kg ha<sup>-1</sup> sulfur, followed closely by those treated with 60 kg ha<sup>-1</sup> sulfur (18.15% oil content and 534 kg ha<sup>-1</sup> oil yield). Conversely, control plots exhibited the lowest oil content (16.90%) and oil yield (448 kg ha<sup>-1</sup>). Application at the flowering stage (R1)

*Table 4 Yield and yield related attributes of soybean under different sulfur levels applied at various application stages.*

Application stages (APP)	Harvest index (%)	Oil content (%)	Oil yield (kg ha <sup>-1</sup> )	Protein content (%)
At sowing stage	33.95	16.67 c	462 c	29.15 b
At V3 stage	34.50	17.50 b	520 b	29.96 b
At R1 stage	35.17	19.54 a	556 a	34.28 a
LSD <sub>(0.05)</sub>	ns	0.71	11	1.07
Sulfur levels (S)				
Control	32.92 c	16.90 c	448 d	29.77 c
20 kg ha <sup>-1</sup>	33.79 bc	17.60 bc	496 c	30.80 bc
40 kg ha <sup>-1</sup>	34.67 b	18.95 a	571 a	31.56 ab
60 kg ha <sup>-1</sup>	36.78 a	18.15 ab	534 b	32.39 a
LSD <sub>(0.05)</sub>	1.47	0.82	13	1.23
APP×S	ns	ns	ns	**

The use of different alphabetic labels within the same groups indicates statistically significant differences at a 5% probability level.

### *Association between different attributes under different sulfur levels at various application stages in soybean*



Scatter plots explained the association between studied variable when evaluated

against different sulfur levels applied at various growth stages. It was found out that plant height was positively associated with biological yield and leaves number plant<sup>-1</sup> with a Pearson's correlation coefficient (R<sup>2</sup>) of 0.56 and 0.80, respectively (Figure 1). The increase in biological yield and leaves number was prominent when the height of plants improved. The association of grain yield with pods plants<sup>-1</sup> and TGW was not significant ( $P = 0.07$  and  $0.51$ , respectively).



However, the correlation was observed to be positive between grain yield and oil yield ( $R^2 = 0.66$ ) that showed that improvement in oil yield was notable when grain yield of soybean was increased (Figure 2). Moreover,

positive and significantly higher correlation was observed between oil and protein content ( $R^2 = 0.92$ ) that revealed synergistic effect between oil and protein content of soybean (Figure 3



).

Figure 1: Scatter plots for plant height VS biological yield and leaves plant-1 under different sulfur levels applied at various application stages




Figure 2: Scatter plots for Grain yield VS pods plant-1, TGW and oil yield under different sulfur levels applied at various application stages.




Figure 3: Scatter plots for Protein content VS oil content under different sulfur levels applied at various application stages

### Economic analysis

In our finding economic analysis indicated that the net profit from harvest of the soybean was more than the cost on its cultivation. Data regarding economic study of the production from soybean are presented in Table 5. It presents the total experimental field area ( $432 \text{ m}^2$ ), which included labor cost (Rs.700  $\text{day}^{-1}$ ), seed cost (Rs.250 @ Rs.80  $\text{kg}^{-1}$ ), elemental sulfur cost (Rs.180), straw cost (Rs.5  $\text{kg}^{-1}$ ), urea cost (Rs.72  $\text{kg}^{-1}$ ) and cost of SSP (Rs.36  $\text{kg}^{-1}$ ). Maximum net income (206334 /-) was obtained when 40  $\text{kg}$  sulfur  $\text{ha}^{-1}$  was introduced at reproductive stage as compared to control (186494 /-).

Table 5: Economic analysis of soybean as affected by sulfur levels and application stages

Production Technology	Production (kg ha <sup>-1</sup> )		Output Value (Rs. ha <sup>-1</sup> )			Total Charges (Rs. ha <sup>-1</sup> )						Net income	B C R
	Grain yield	Straw yield	Grain value	Straw value	Gross income	N cost	P cost	Sulfur cost	Labor cost	Seed cost	Total cost		
0	2547	7745	178290	38725	217015	596	155	0	5600	20000	26351	190664	6.9
20	26450	7830	185150	39150	224300	596	155	4320	5600	20000	30671	193629	6.2
40	28827	7917	201740	39585	241325	596	155	8640	5600	20000	34991	206334	5.5
60	27395	7855	191730	39275	231005	596	155	12960	5600	20000	39311	191694	4.6
Application Stages													
Sowing	25883	7633	181160	38165	219325	596	155	6480	5600	20000	32831	186494	5.6
Vegetative	27174	7884	190190	39420	229610	596	155	6480	5600	20000	32831	196779	4.9
Reproductive	28062	7992	196420	39960	236380	596	155	6480	5600	20000	32831	203549	6.1

Whereas,

Price of soybean seed = Rs. 250 kg<sup>-1</sup>

Price of straw = Rs. 5 kg<sup>-1</sup>

Price of grain = Rs. 70 kg<sup>-1</sup>

Cost of sulfur = Rs. 180 kg<sup>-1</sup>

Cost of urea (N) = Rs. 72 kg<sup>-1</sup>

Cost of ssp. (P) = Rs. 36 kg<sup>-1</sup>

Labor charges = Rs. 700 day<sup>-1</sup>

## DISCUSSION

The analyzed results showed that sulfur levels and application stages notably impacted the phenological development, specifically the duration taken for flowering and physiological maturity in soybeans. Higher sulfur dose significantly prolonged the days to flowering in soybean i.e. supplementing 40 kg ha<sup>-1</sup> sulfur caused delay in flowering in comparison with control plots. This delay in flowering associated with higher sulfur levels could be attributed to sulfur role in augmenting chlorophyll content and enhancing photosynthetic processes, consequently fostering improved and prolonged vegetative growth in the crop. A study by Shipley et al. (2012) noted that increased sulfur levels led to delay in flowering in soybean, particularly observing delayed flowering occurrences at the 40 kg S ha<sup>-1</sup> level. Application stages (APP) of sulfur significantly affect days to flowering and maturity. Sulfur applied at vegetative stage (V3) delayed flowering. While minimum was recorded at sowing stage. Similarly, delayed maturity was seen at 60 kg S ha<sup>-1</sup> as compared to control plot. Sulfur applied at flowering stage (R1) had delayed maturity as compared to sowing stage in soybean. According to Haruna et al. (2011) late flower formation in soybean might be due to high nitrogen and sulfur application which produce vigorous plant growth and prolonged vegetative growth stage resulted in delay flowering and physiological maturity.

Plants subjected to higher sulfur levels, particularly at 40 kg ha<sup>-1</sup>, displayed taller plants with maximum number of leaves. This enhancement linked to increased sulfur levels could be attributed to sulfur role as a constituent in plant tissue, significantly influencing soybean height, as indicated by Shahid et al. (2009). Conversely, the decrease observed in these parameters at lower sulfur application levels might be due to the stunted growth and limited development of plants under sulfur deficiency. Consistent with these findings, studies by Varun et al. (2011) and Burkitbayev et al. (2021) noted that sulfur led to enhanced plant height and

increased leaf count per plant. Their rationale centered on the capacity of sulfur to alter soil physicochemical properties, thereby enhancing nutrient availability and fostering plant growth and development. This, in turn, facilitated nutrient translocation towards generative organs, positively influencing photosynthesis, and ultimately contributing to augmented yields and structural yield elements. Moreover, Begum et al. (2015) noted that the tallest plants and more branches were evident in the high sulfur treatment, surpassing the lower treatment groups. Concerning sulfur application stages, maximal plant height and leaf count were observed when sulfur was applied during the vegetative stage (V3). Conversely, the lowest plant height and leaf count were observed when sulfur was applied at the sowing stage. This disparity might be attributed to the duration of the vegetative period being shorter compared to other stages, particularly given that most soybean cultivars are categorized as short-day plants. This aligns with Kumar et al. (2017) affirming the positive impacts of fertilizer application during the vegetative phase on the physiological aspects of soybean cultivation, particularly on plant height and leaf count per plant.

Among yield components of soybeans, specifically rise in pod count per plant was evident with an increasing application of sulfur, especially up to 40 kg S ha<sup>-1</sup>, alongside a simultaneous increase in pod length. This improvement can be linked to the essential role of sulfur in diverse plant growth and developmental processes. Sulfur, a vital element in amino acids containing sulfur, may have played a role in fostering the transition of tissue from somatic to reproductive states, stimulating meristematic activity, and contributing to overall plant development, which resulted in increased pod quantity and size. These observations align with the findings of Chaubey et al. (2000), who noted significant enhancements in pod quantity, length with sulfur. TGW weight was positively influenced by higher levels of sulfur when applied at R1 stage of growth. This might be due to sulfur that enhanced nutritional conditions,

leading to improved carbohydrate metabolism and aiding in the transfer of photosynthates to seeds, thereby resulting in larger grains. This observation is corroborated by research indicating a correlation between sulfur treatment and increased seed weight as stated by Sharifi et al. (2012). Similarly, applying sulfur at a rate of 40 kg S ha<sup>-1</sup> during the reproductive stage (R1) produced higher grain yield compared to sowing stage application. Dhaker et al. (2010) also documented a significant rise in soybean seed yield with 40 kg S ha<sup>-1</sup> application. Likewise, Casteel et al. (2019) reported a substantial increase in soybean grain yield in response to sulfur application in their study on soybean production.

Commencement of sulfur and the timing of sulfur application significantly influenced biomass yield and harvest indices of soybeans. Specifically, sulfur at a rate of 40 kg ha<sup>-1</sup> during the reproductive stage (R1) notably increased the biological yield. Moreover, applying 60 kg ha<sup>-1</sup> sulfur during the reproductive stage led to a higher harvest index. Shah et al. (2013) reported that escalating sulfur rates substantially enhanced the biological yield by stimulating chlorophyll content and protein synthesis, promoting increased photosynthesis and ultimately elevating biomass production. It is crucial to highlight that the adequacy of soil fertilization and subsequent plant nutrition plays a major role in intensifying agricultural production, as emphasized by Diacono and Montemurro (2010).

Elevating dose of sulfur from 0 to 40 kg ha<sup>-1</sup> resulted in a substantial increase in both the oil content and oil yield. However, further increasing the sulfur level resulted in a reduction in both oil content and yield. This change could be linked to sulfur's role in synthesizing fatty acids and improving protein quality by aiding in the production of particular amino acids (Konyak et al., 2016). When sulfur was applied during the flowering stage (R1), it resulted in higher oil content and yield compared to the application at the sowing stage. This difference could be due to increased metabolic activity in soybean crops

during the flowering stage, leading to an enhanced production of secondary metabolites. Additionally, a preference for allocating cysteine towards glutathione and protein synthesis might contribute to these differences (D'Hooghe et al., 2013). In last, protein content was soybean differed significantly by different levels of sulfur. Studies have shown that sulfur-containing agrochemicals, particularly in the form of powdery and solute applications, lead to higher growth, yield, and protein content in soybeans compared to control conditions (Głowacka et al., 2023). Application of sulfur fertilizer has been found to positively impact protein concentrations and oil content in soybeans, ultimately enhancing yield and nutritional quality as also claimed by Mahato, (2023).

### **Conclusion**

In conclusion, the study underscores the significant effects of sulfur levels and application stages on soybean growth and yield components. The most beneficial results were achieved with sulfur applied at a rate of 40 kg ha<sup>-1</sup>, which enhanced plant height, leaf number, pod size, and both biological and grain yields. Moreover, this sulfur level improved oil and protein content, thus boosting overall seed quality. For farmers, these findings imply that incorporating sulfur into their agronomic practices can lead to substantial improvements in soybean productivity. Specifically, applying sulfur at the flowering stage (R1) is recommended, as it effectively maximizes seed, oil yields and economically friendly. Adopting this practice could enhance crop performance and profitability, making it a valuable practice in soybean cultivation in the region.

### **Conflict of interest**

All the authors who contributed in the manuscript declared no conflict of interest.

### **Authors' contributions**

The contributions of all the authors are appreciated and acknowledged.



## References

- Ahmad, A., Abrol, Y.P. and Abdin, M.Z., 2011. Effect of split application of sulfur and nitrogen on growth and yield attributes of Brassica genotypes differing in time of flowering. *Canadian Journal of Plant Science*, 79, 175-180.
- Ahmad, G., Jan, A., Arif, M., Jan, M.T. and Khattak, R.A., 2007. Influence of nitrogen and sulfur fertilization on quality of canola (*Brassica napus* L.) under rainfed conditions. *Journal of Zhejiang University Science B*, 8,731-737.
- Ahmad, J., Ahmad, F., Iqbal, S., Shah, S.M.A., Ali, M., Abbas, M.W., Nawaz, H., Mehmood, Z., Ali, B. and Ali, S., 2018. Growth and oil yield of sesame as influenced by sulfur and nitrogen. *Research in Agriculture, Livestock, and Fisheries*, 3, 147-150.
- Begum, A., Roy, M., Roy, D.C., Sujit, A., 2015. Performance of soybean in different levels and time of sulfur application in Indo-Gangetic plains of eastern India. *Journal of Experimental Biology and Agricultural Sciences*, 6, 929-935.
- Berger, K.C., Truog, E., 2010. Sulfur determination in soils and plants. *Industrial & Engineering Chemistry Analytical Edition*, 11, 540-545.
- Burkitbayev, M., Bachilova, N., Kurmanbayeva, M., Tolenova, K., Yerezhopova, N., Zhumagul, M., Mamurova, A., Turysbek, B., Demeu, G., 2021. Effect of sulfur containing agrochemicals on growth, yield, and protein content of soybeans (*Glycine max* L.). *Saudi Journal of Biological Sciences*, 28, 891-900.
- Casteel, S.N., Chien, S.H., Gearhart, M.M., 2019. Field evaluation of ammonium sulfate versus two fertilizer products containing ammonium sulfate and elemental sulfur on soybeans. *Communications in Soil Science and Plant Analysis*, 50, 1-7.
- Chaubey, A.K., Singh, S.B., Kaushik, M.K., 2000. Response of groundnut (*Arachis hypogaea* L.) to sources and level of sulfur fertilizer in mid-western plains of Uttar Pradesh. *Indian Journal of Agronomy*, 45, 166-169.
- D'Hooghe, P., Escamez, S., Trouverie, J., Avice, J.C., 2013. Sulfur limitation provokes physiological and leaf proteome changes in oilseed rape that lead to perturbation of sulfur, carbon, and oxidative metabolisms. *BMC Plant Biology*, 13, 1-15.
- Dhaker, S.C., Mundra, S.L., Nepalia, V., 2010. Effect of weed management and sulfur nutrition on productivity of soybean (*Glycine max* L.). *Indian Journal of Weed Science*, 42, pp.232-234.
- Diacono, M., Montemurro, F., 2010. Long-term effects of organic amendments on soil fertility. A review. *Agronomy for Sustainable Development*, 30, 401-422.
- Dubey, S.K., Billore, S.D., 1995. Effect of level and sources of sulfur on symbiotic and biometrical parameters of soybean (*Glycine max* L.). *Indian Journal of Agricultural Sciences*, 65, 140-144.
- Głowacka, A., Jariene, E., Flis-Olszewska, E., Kiełtyka-Dadasiewicz, A., 2023. The effect of nitrogen and sulphur application on soybean productivity traits in temperate climates conditions. *Agronomy*, 3,780.
- Haruna, L.M., 2011. Dry matter partitioning and grain yield potential in sesame (*Sesamum*

- Indicum* L.) as influenced by poultry manure, nitrogen and phosphorus at Samaru. Nigeria Journal of Agricultural Technology, 7, 1571-1577.
- Hossain, M.S., Alam, M.N., Mandal, S., Rashid, M.M., Morshed, M.N., 2019. Effect of sulfur and zinc on the yield and oil content of soybean. Bangladesh Journal of Environmental Science, 36, 39-42.
- Jan, M., Shah, T.P., Hollington, P.A., Khan, M.J., Sohail, Q., 2009. Agriculture research design and analysis a monograph NWFP Agriculture University Peshawar, Pakistan.
- Kahraman, A., 2017. Nutritional value and foliar fertilization in soybean. Journal of Elementology, 22, 55-66.
- Karami, S., Sanavy, S.A.M., Sanam, S., Keshavarz, H., 2016. Effect of foliar Zinc application on yield, physiological traits and seed vigor of two soybean cultivars under water deficit. Notulae Scientia Biologicae, 8, 181-191.
- Kjeldahl, C., 1883. A new method for the determination of nitrogen in organic matter. Anal of Chemistry, 22, 366.
- Kumar, S., Wani, J.A., Lone, B.A., Fayaz, A., Singh, P., Qayoom, S., Dar, Z.K., Ahmed, N., 2017. Effect of phosphorus and sulfur on nutrient and amino acids content of soybean (*Glycine max* L.) under 'Alfisols'. Journal of Experimental Agriculture International, 16, 1-7.
- Mahato, G.R., 2023. Effect of Foliar and Preplant Sulfur Application on Soybean Nutrient Uptake, Yield, and Seed Protein and Oil Content. South Dakota State University.
- Mahmood, B., Abba, A., Daliri, M.S., Namdari, M., 2013. The evaluation of different values of phosphorus and sulfur application in yield, yield components and seed quality characteristics of soybean (*Glycine max* L.). Advances in Environmental Biology, 7, 170-176.
- Ministry for Food Agriculture and Livestock (MNFS&R), 2014. Agriculture Statistics of Pakistan. Government of Pakistan, Economic Wing, Islamabad.
- Mukhtar, M., Klosterman, D., Benin, V., Morgan, A., 2016. Synthesis of a phosphorus-based epoxy reactive flame-retardant analog to diglycidyl ether of bisphenol A (DGEBA) and its behavior as a matrix in a carbon fiber composite. Polymer Degradation and Stability, 205, 110-144.
- Paul, S.K., Khatun, M.M., Sarkar, R., 2019. Effect of sulfur on the seed yield and oil content of soybean (*Glycine max* L.). Journal of Bangladesh Agricultural University, 17, 33-38.
- Seervi, D.P., Choyal, K.S., Seervi, K.S., 2018. The effect of micronutrients applied as foliar spray, on yield & yield attributes and oil content of sesame crop (*Sesamum indicum* L.). Journal of Pharmacy and Phytochemistry, 7, 1402-1404.
- Shah, M., Manaf, A., Hussain, A., Farooq, M., Zafar-Ul-Hye, M., 2013. Sulfur fertilization improves the sesame productivity and economic returns under rainfed conditions. International Journal of Agriculture and Biology, 15, 1301-1306.
- Shahid, M.Q., Saleem, M.F., Khan, H.Z., Anjum, S.A., 2009. Performance of soybean (*Glycine max* L.) under different sulfur levels and inoculation. Pakistan Journal of Agricultural Sciences, 46, 237-241.

- Shilpi, S., Islam, M.N., Sutradhar, G.N.C., Husan, A., Akter, F., 2012. Effect of nitrogen and sulfur on the growth and yield of sesame. *International Journal of Bioresource Stress Management*, 3, 108-111.
- Shivakumar, B.G., Gangaiah, B., Ahlawat, I.P.S., 2009. Effect of sulfur nutrition on performance of soybean-wheat cropping sequence in northern plain zone of India. *Journal of Oilseed Research*, 26, 385-387.
- Varun, B.K., Patel, B.A., Panchal, H.D.B., Dalwadi, M.R., Patel, J.C., 2011. Effect of different sources and levels of sulfur on yield and quality of soybean under inceptisol of middle Gujarat. *Asian Journal of Soil Science*, 6, 85-87.
- Vora, V.D., Hirpara, D.S., Vekaria, P.D., Sutaria, G.S., Vala, F.G., 2018. Effect of sulfur management on yield, nutrient uptake by soybean and post-harvest soil fertility under rainfed condition. *Journal of Pharmacy and Phytochemistry*, 7, 65-69.
- Zhao, Y., Bi, D., Zhao, Q., Liu, C., Hu, Z., 2006. Physiological and ecological effects of sulfur fertilization on soybean. *Journal of Applied Ecology*, 17, 2376-2380.